JOURNAL OF PURE AND APPLIED ALGEBRA
VOLUME 139, NUMBERS 1–3, 17 JUNE 1999

Effective Methods in Algebraic Geometry

CONTENTS

Editorial 1

P.H. Berman and M.F. Singer
Calculating the Galois group of $L_1(L_2(y))=0$, L_1, L_2 completely reducible operators 3

W.A. de Graaf
Using cartan subalgebras to calculate nilradicals and Levi subalgebras of Lie algebras 25

N. Dutertre
An algebraic formula for the Euler characteristic of some semi-algebraic sets 41

J.-C. Faugére
A new efficient algorithm for computing Gröbner bases (F_4) 61

R. González-Díaz and P. Real
A combinatorial method for computing Steenrod squares 89

M. van Hoeij
Finite singularities and hypergeometric solutions of linear recurrence equations 109

S. Hoşten and R.R. Thomas
Standard pairs and group relaxations in integer programming 133

G. Ivanyos
Finding the radical of matrix algebras using Fitting decompositions 159

M.G. Marinari and L. Ramella
Some properties of Borel ideals 183

T. Oaku and N. Takayama
An algorithm for de Rham cohomology groups of the complement of an affine variety via *D*-module computation 201

D. Richardson
Elimination of infinitesimal quantifiers 235

D. Rupprecht
An algorithm for computing certified approximate GCD of *n* univariate polynomials 255

M. Stillman and H. Tsai
Using SAGBI bases to compute invariants 285

U. Walther
Algorithmic computation of local cohomology modules and the local cohomological dimension of algebraic varieties 303

T. Zell
Betti numbers of semi-Pfaffian sets 323

Author Index 339